An Affirmative Action/Equal Opportunity Eq

### Edited by Jill Warren Photocomposition by Barbara J. Velarde

NOTICE: Reproduction of this document requires the written consent of the originator, his successor, or higher authority.

### DISCLAIMER

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.





### LOW-YIELD NUCLEAR EXPLOSION CALCULATIONS: THE 9/22/79 VELA SIGNAL (U)

by

E. M. Jones, R. W. Whitaker, H. G. Horak, and J. W. Kodis

### ABSTRACT (SRD)

DOE b(1)

Normally, independent confirming evidence is available from other VELA satellites and other sources. Unfortunately, such confirming evidence has not been uncovered for the 9/22/79 event.

DOE (1)

In this report we summarize the veia data, discuss classical interpretations, and present a particular model which, we believe, satisfactorily reproduces the Vela signal

DOE b(1)

Thus, our model is consistent with the apparent absence of nuclear debris, the collection of which is required by some analysts for absolute confirmation of an atmospheric detonation.

### I. THE VELA DATA

The VELA satellite and its detectors are described elsewhere (e.g. Horak 1980). For our purposes, the salient features of the detections system are

(:


(4

1 DOE

bu.







DOE B(i)

observed by the VELA satellite. As a first step, we use classic scaling laws to estimate the significant parameters of yield and burst height density; then we discuss computer calculations chosen to provide more precise estimates of the explosion parameters.

Numerous sets of scaling laws exist. Here, we have synthesized four adequate scaling laws from discussions by Zinn et al. (1974) and Sappenfield (1979).

As we shall see, the time of first maximum and the power ratios are inconsistent with the other observations. Solving the puzzle will concern most of the rest of our study.

DOE b(1)

Doe pui


Clearly, conventional scaling laws cannot provide a consistent model for the VELA data.

The values of the observable quantities (t<sub>1-max</sub>, t<sub>2-max</sub>, and P<sub>2-max</sub>P<sub>min</sub>) have large uncertainties.

6(1)



Both Y and p are model independent because they are the basic parameters of any calculation. The other three factors can vary dramatically among different models. Let us now consider these other factors and discuss how they can affect the signal.





significant modification of the signal by the atmospher or by clouds.

Nuclear weapons with significant neutron output can create an obscuring curtain of "smog" (principally NO<sub>2</sub>, HNO<sub>2</sub>, and O<sub>3</sub>). As discussed by Zinn et al. (1974), smog may produce not only a delay in first maximum but also a severe depression of the first pulse by absorption. The minimum and second maximum usually are unaffected by smog because the fireball radius at these times exceeds the distances at which significant neutron deposition occurs.

Perhaps the most dramatic mass effect has been known since the earliest days of atmospheric testing: the apparent yield of a burst at the earth's surface (land or sea) is twice the actual yield. The reason is simply that the surface acts as a nearly perfect reflecting plane. The entire explosion energy acts only in the hemisphere above the surface as if the surface were absent and we were witnessing the spherical expansion of a higher yield explosion. The result is that the radiative/hydrodynamic behavior is identical to that of a free-air burst at twice the actual yield. Among others, Sappenfield (1979) has reviewed the empirical evidence supporting this result.

Cloud cover, atmospheric absorption, and scattering by particulates between the explosion and the satellite can alter the interpretation of the signal. Atmospheric transmission must be considered when deriving radiance at the source from the observed irradiance. However, simple absorption by the atmosphere will not produce dramatic shifts in the timings of events in the signal nor, to first order, will it depress one part of the signal relative to another. However, cloud cover can introduce time smearing caused by photon scattering. Multiple photon scatterings can delay first maximum but should also depress the signal in both peaks. The observed irradiance levels are high enough that the signal could not have suffered much absorption nor scattering beyond that expected for relatively clear maritime air. We do not believe that the data supports the contention that there was any

The hydrodynamic behavior and the power-time curve of this case are illustrated in Figs. 3 and 4, respectively.

P(3)





output can

atmosphere

DOE b(3)



UNCLASSIFIED



Dοξ b(3)

The rest of this report discusses the effects of mass on the complete optical signal.

### III. RADFLO CALCULATIONS

For our present purposes, we have chosen to do calculations in which the nuclear explosive is modeled as a sphere of high-density air at uniform density. The "air bomb" comprises the 10 innermost cells in the calculational grid

In each calculation, a certain fraction of the explosion energy is deposited as internal energy at t = 0 in the central cell.

What matters here is the kinetic energy content of the debris, which is largely insensitive to the material composition.

The principal results are tabulated in Tables II and III.

The effect of M and  $\rho$  on the times of minimum and of the two maximums are shown in Figs. 7-9. The choice of a time value from a calculation or from data is subjective.

DOÉ 5(3)

DOE b(3)

D2€ D2€

POS)

WEUNCLASSIFIEM BELLAND

DOE W3)

D56



UNCLASSIFIED

ONOTHER TIT

DOE 6(3)



UNCLASSIFIED

DOE P(3)

Det .

Dog )

Dok 3

(12)

The agree-

ment is satisfactory.

After the shock emerges from the bomb and begins to sweep up ambient air, the power-time curve shows dramatic oscillations produced by the entrainment of individual calculational cells. When the shock first encounters an ambient cell, the air temperature in the cell rises, causing the opacity to rise and thereby decreasing fireball brightness. Eventually, the newly entrained cell is heated enough that it starts radiating, and the fireball brightness increases until the cycle is repeated with the entrainment of the next cell outward. Interpretation of the calculated power levels during the first pulse is uncertain. We have chosen to show the power levels averaged over each cycle as

$$\bar{P} = \frac{1}{\delta t} \int_{\text{cycle}} P \, dt \ . \tag{13}$$

The calculated amplitude of the bump at high mass is well below the data uncertainties.

At later times, physical mechanisms responsible for producing the power-time curve variations are well understood. The interested reader may consult Zinn (1973), Zinn et al. (1974), Brode (1968), or Glasstone (1964) for discussions of phenomena at t<sub>min</sub> and beyond.

We have plotted our best interpretation of the VELA data to produce the power outputs in Fig. 12.

## V. PERTINENT DATA FROM ATMOSPHERIC NUCLEAR TESTS

We have constructed what we believe to be a plausible model for a low-yield nuclear explosion that could have

Doé bos



VIn this section, we examine the available data for t<sub>1-m</sub> from atmospheric nuclear tests.

The available t<sub>1-max</sub> data are plotted as a function of yield in Fig. 13. We note that for many events of the US atmospheric test program, data pertinent to the first maximum are not available.

However, the fit after about 1 ms

is quite satisfactory.

D)É

\*Personal communication to authors.

Dog (3)




Doé b(1) p(3)

DOE DOS

GEORGE was detonated atop a tower 61 m above Eniwetok on May 9, 1951

706. 613

phenomenology are of interest. First, the fireball expansion was very asymmetric during the early phases; second, independent determination of the yield by hydrodynamic and radiochemical techniques were widely discrepant.



360

DOE 100

Dot b(3)

The other part of the fireball may have grown radiatively. Clearly, calculational studies in two or three dimensions will be required to provide a convincing explanation.

At later times, the expansion becomes more symmetric and, at minimum, this fireball has prominent spots.

206 Spers.

The hydrodynamic yield is determined by comparing the radius-time data with the expectations of classic blast wave theory. In particular, during the period immediately preceding minimum, the quality

is virtually constant. In Fig. 16, we show the  $\phi$  histories obtained from four RADFLO calculations compared



VI. CAVEATS

with the GEORGE data. These data were derived by visually estimating a "best-circle" fit to the fireball shape.

Nonetheless, we recognize that we have made certain assumptions that, if proved wrong, may alter our conclusions.

(1)

(2) We have used dense air to model the weapon vapor.

b(3)

DOE 6(3)



(3)

have been unable to secure experimental confirmation of a t<sub>1-max</sub> delay caused by mass.

DOE (3)

(4)

The

possibility remains that other models can produce good fits to the data. Proponents of such models must demonstrate plausibility.

VII. SUMMARY

# DOÉ (3)

\*Private communication to authors.

#### **ACKNOWLEDGMENTS**

We gratefully acknowledge valuable discussion with H. Hoerlin, J. Zinn, and J. Malik of Los Alamos National Laboratory and G. Mauth of Sandia National Laboratories.

#### REFERENCES

Brode, H. L., "Numerical Solutions of Spherical Blast Waves," J. App. Phys. 26, 766 (1955).

Brode, H. L., "Review of Nuclear Weapons Effects," Ann. Rev. Nuc. Sci. 18, 153 (1968).

Colvin, J. D. and N. P. Gow, "Fireball Spots: A Comprehensive Study of the Photographic Data (U)," EG&G Technical Report EGG 1183-5086 (SFRD) (August 1978).

Glasstone, S. (Ed), The Effects of Nuclear Weapons - Revised Edition (U) (Washington: U.S. Atomic Energy Commission 1964).

Horak, H. G., "Vela Event Alert 747 (U)," Los Alamos Scientific Laboratory Report LA-8364-MS (SRD) (May 1980).

Horak, H. G. and J. W. Kodis, "RADFLO: A User's Manual (U)," Los Alamos National Laboratory Report LA-9245-M (U) submitted for publication.

Sappenfield, D. S., "The Surface Burst Correction Factor for Bhangmeter Scaling (U)," Air Force Technical Applications Center Topical Report AFTACTR-79-12 (SRD) (March 1979).

Sedov, L. I., Similarity and Dimensional Methods in Mechanics (New York: Academic Press, 1959).

Taylor, G. I., "The Formation of a Blast Wave by a Very Intense Explosion, I. Theoretical Discussion," Proc. Roy. Soc. A201, 159 (1950).

Zinn, J., "A Finite Difference Scheme for Time-Dependent Spherical Radiation Hydrodynamics Problems," J. Comp. Phys. 13, 569 (1973).

Zinn, J., J. W. Kodis, and C. D. Sutherland, "Status Report on Fireball Radiation Hydrodynamics Computations (U)," Los Alamos Scientific Laboratory Report LA-5591-SR (SFRD) (May 1974).



### Standard Distribution

	Copy No.		Copy No.
US DOE, Headquarters Library, Reports Section, Washington, DC	1-3	Commander, Naval Sea Systems Command, Washington, DC	13
Manager, ALO, Albuquerque, New Mexico	4	Director, Air Force Weapons Laboratory,	14-16
Lawrence Livermore National Laboratory, Livermore, California	5-6	Kirtland AFB, New Mexico	
Sandia National Laboratories, Albuquerque, New Mexico	7	US DOE, Technical Information Center, Oak Ridge, Tennessee	. 17
Military Liaison Committee, Washington, DC	8	Attn: Weapon Data Index	
Headquarters, Defense Nuclear Agency, Washington, DC		Los Alamos National Laboratory Report Library	18-64
Defense Nuclear Agency Field Command, Kirtland AFB, New Mexico	11-12		

### Special Distribution

	opeciai 1	Sistibution	
	Copy No.		Copy No.
Central Intelligence Agency	65-82	Defense Nuclear Agency, Headquarters	94-97
Washington, DC		Department of Defense	
Attn: Chief, OSWR/NED		Washington, DC	
For: Frank P. Baranowski, JAEIC		Director	
Donald M. Brasted, C/NED (2 cys.)		Attn: PAAD-3	
LT. GEN. Harold C. Donnelly, JAEIC		For: C. Fitz	
LTC. James Fargo, DAMI-FIT		L. Wittwer	
Kenneth Gimbert, NSA/AOS		G. Sevin	
LT. COL. Houston Hawkins,		C. Knowles	
DJA(DT-1E)			
Dr. John S. Ingley, OSWR		Air Force Technical Applications Center	98
Dr. Gerald W. Johnson		Patrick Air Force Base, Florida	
Max A. Koontz, NISC		AFTAC/TD	
Robert Morrison, ACDA		Attn: Dr. Gerard M. Leies	
Dr. S. A. Newton, Jr., JAEIC			
Louis H. Roddis, Jr., JAEIC		Air Force Weapons Laboratory	99-100
Robert Rubinstein, DOE		Kirtland Air Force Base, New Mexico	
Jeffry Siegel, JAEIC		Attn: SUL, Mrs. Georgiana Hillyer	
Dr. Gerald F. Tape, JAEIC		For: MAJ. Ganong	
CAPT. Raymond Vitkus, AF/INET			
VADM Steven A. White		Union Carbide Corporation Y-12 Plant	. 101
		Nuclear Division	
Lawrence Livermore National Laboratory	83-86	Oak Ridge, Tennessee	
Livermore, California		Attn: Y-12 Plant Records Department	
Attn: C. Alonso		For: Paul R. Vanstrum	
R. Klein			
Dr. Roger E. Batzel		Harold M. Agnew, Consultant	102
Dr. Edward Teller		c/o General Atomic Company	
		La Jolla, California	
Sandia National Laboratories	87-90		
Albuquerque, New Mexico		Walter G. Chestnut, Consultant	103
Attn: Mail Services Section		Menlo Park, California	
For: Glenn Fowler			
A. Chabai		W. E. Ogle, Consultant	104
G. Mauth		Energy Systems, Inc.	
D. Thornborough		Anchorage, Alaska	
		Attn: W. E. Ogle	
EG&G, Inc.	91-93		
Los Alamos, New Mexico			
Attn: Document Control			
For: D. Wright			
	,		

J. Colvin L. Rauber

